

REMOVAL OF CYSTS IN THE THYROID, OVARIES, AND PANCREAS

FOREWORD. *This process has been successfully tested for several years.* Before reading about any of the treatments, you should learn about TENS (Transcutaneous Electrical Nerve Stimulator), which is equivalent to medicine, and see its characteristics by reading the corresponding file contained in this same package. You can purchase it online, just make sure that the "Stimulation Modes" include the option: "Normal or Constant." If this treatment does not cure you, you should see your treating physician.

TREATMENT PROCESS

The treatment is designed so that the current eliminates all types of cysts in the indicated areas, as well as others in different parts of the body. The treatment always follows its basic principles, one of which states that current must be passed through the area of interest and, therefore, the position in which the pads should be placed must be designed.

TREATMENT PARAMETERS

TIME. *This is indicated in the treatment description.*

CURRENT INTENSITY. *Very gently increase the corresponding TENS knob until you feel a slight current, without pain, and leave it at that level for the time indicated in the treatment description; this should be the intensity used for all treatments and conditions.*

TREATMENT OF THE THYROID GLAND.

This is a gland located in the throat, and its electrical treatment must follow a proper medical diagnosis of the condition, which can have various causes: nodules, cysts, cancer, etc.



Figure 1. Image of the thyroid gland.

1. Place pads on both sides of the throat (position 1 marked with red pads in Figure 2). Stimulate for 10 minutes at the recommended intensity.





Figure 2. Placement of pads for thyroid stimulation.

2. Placement of pads between the tongue and the base of the



neck.

Figure 3. Stimulation of the inside of the throat toward the thyroid gland.

Figure 3 illustrates how the pads should be placed; treat for 15 minutes in this position at the recommended intensity.

TREATMENT OF OVARIAN CYSTS.

1. Place one pad at the entrance to the vagina (pad 1) and another on the abdomen, in the indicated position (pad 2 on the right side of the abdomen); stimulate




Figure 4. Treatment of ovarian cysts.

Continue for 10 minutes. Then move the pad on the abdomen to the symmetrical left side (position 3) and stimulate there for 10 minutes. Pad 1 does not move during the treatment.

PANCREATIC CYSTS


This disease is similar to that described above for the thyroid and ovaries. In this case, lateral treatment is performed by placing pads on both sides of the chest and abdomen and another pad on the front.

SIDE TREATMENT


As shown in Figure 5(a), the left pad is placed between ribs 3 and 4, counting from bottom to top, and in the rear half of the abdomen, or closer to the back. The second pad goes on the right side, below the last rib, as shown in Figure 5(a); with the pads in this position, stimulate for 15 minutes at the recommended intensity in order to cover the entire length of the pancreas.

FRONTAL TREATMENT

To reinforce the treatment, it is recommended to stimulate the abdomen frontally (from front to back), placing one pad on the right side of the spine and below the ribs on the back, as shown in Figure 5(b), which shows part of the pancreas in yellow. The second pad goes on the front at the same height as the back pad. Stimulate for 10 minutes in this position.



(a)



(b)

Figure 5(a). Position of the pancreas and pads in frontal view; (b) View of the pancreas from the back.