

TREATMENT OF GASTROINTESTINAL INFECTIONS

FOREWORD. *This process has been successfully tested for several years and its effectiveness has also been verified by the global community.* Before reading about any of the treatments, you should learn about TENS (Transcutaneous Electrical Nerve Stimulator), which is equivalent to medicine, and see its characteristics by reading the corresponding file contained in this same package. You can purchase it online, just make sure that the "Stimulation Modes" include the option: "Normal or Constant." If this treatment does not cure you, you should see your treating physician.


Treatment

For example, if the person has nausea or vomiting, it is very likely that these infectious agents are in the stomach; if, on the other hand, they only have diarrhea, with or without blood, and a fever, the infection is mainly located in some part of the intestine; and if they have both, the disease affects both the stomach and the intestine.

The application of electric current in the treatment of diseases has always impressed me in a very pleasant way, healing almost immediately, as is the case with active herpes, in just 10 minutes with 100% effectiveness.



(a)



(b)

**Figure 1. Positions of the stomach (a)
and intestine (b)**

.

TREATMENT

Figure 1 shows the physical positions of both the stomach and the intestine, which also makes clear where the pads should be placed to eliminate the microbes causing the disease. Remember that the current always takes the path of least resistance between the two pads; and because the body is mostly water, it will tend to travel almost in a straight line between them once inside the body. For this reason, a natural path would be to follow the same path as food: it enters through the mouth and finally exits through the anus. This is the most logical or natural path and can be used as the final part of the treatment for 5 minutes. Obviously, the pad in the anus must be disposed of safely by sticking it to a double Kleenex and then placing it in a plastic bag.

TREATMENT PARAMETERS.

Time. "In all positions, the treatment duration should be 5 minutes."

Intensity. The intensity of the current should be increased until the person feels it, but without causing pain, and left at that level.

Frequency. "The TENS frequency should always be set to the maximum setting of the device, which is normally 150 or 200 hertz, and saved so that it always operates in this way."

STOMACH TREATMENT

PHASE 1 (fig. 2a). From the mouth to the lower abdomen, above the mons pubis or hairy part of the abdomen; in this position, it travels from the mouth, pharynx, larynx, esophagus, part of the stomach, and practically the entire intestine vertically. The current does not follow the shape of the intestine, traveling through all its twists and turns to the anus.

PHASE 2. This focuses mainly on the stomach; from the beginning of the stomach, below the sternum bone where the first pad is placed, and on the left side of the chest below the ribs, about 10 centimeters away from the first, the second pad is placed to cross the stomach in that direction (fig. 2b).



Figure 2. Stomach treatment: One pad on the tongue and the other below the navel.

INTESTINAL TREATMENT

In this case, the design should cover the entire abdomen where the large and small intestines are located in order to eliminate the bacteria or parasites that infect it. That is why in the design in Figure 3, it is swept both vertically and horizontally. In all cases, common sense shapes the design of the pad placement. ***The treatment should last 5 minutes in each position.***

A natural design and suggestion, as mentioned at the beginning, is to place one pad on the tongue and the other on the anus. The sensation of the current in the anus is no more painful than in any other part of the body, and the current sweeps through the space between both ends of the gastrointestinal tract ***for 10 minutes.*** In my opinion, this is a good way to end the treatment, which also increases the likelihood of sweeping through all infected areas.




Figure 3. Treatment of the intestine; the current flows from one pad to the other, as indicated by the different positions. Position 1 is fixed in the anus and does not move. Positions 2, 3, 4, 5, 6, and 7 sweep across the abdomen, irradiating it from the anus.